

AD-A055 829

PENNSYLVANIA STATE UNIV UNIVERSITY PARK COLL OF BUSIN--ETC F/G 12/1  
STOCHASTIC DOMINANCE AND MOMENTS OF DISTRIBUTIONS.(U)

JUN 78 P C FISHBURN

N00014-75-C-0857

UNCLASSIFIED

TR-29

NL

| OF |  
ADA  
055829



END  
DATE  
FILMED  
8 -78  
DDC

UNCLASSIFIED

FOR FURTHER TRAN

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER Technical Report, No. 29	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) STOCHASTIC DOMINANCE AND MOMENTS OF DISTRIBUTIONS.		5. TYPE OF REPORT & PERIOD COVERED Interim
6. AUTHOR(s) Peter C. Fishburn		6. PERFORMING ORG. REPORT NUMBER
7. PERFORMING ORGANIZATION NAME AND ADDRESS The Pennsylvania State University University Park, Pennsylvania 16802		8. CONTRACT OR GRANT NUMBER(s) N00014-75-C-0857
9. CONTROLLING OFFICE NAME AND ADDRESS (code 613B: SJW) Office of Naval Research, Dept. of the Navy 800 N. Quincy Street Arlington, Virginia 22217		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 121407
11. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) 1214p. 14 TR-29		12. REPORT DATE June 1978
		13. NUMBER OF PAGES 12
		14. SECURITY CLASS. (of this report) Unclassified
15. DISTRIBUTION STATEMENT (of this Report) Distribution of this document is unlimited.		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
17. SUPPLEMENTARY NOTES		
18. KEY WORDS (Continue on reverse side if necessary and identify by block number) Stochastic dominance Moments of distributions		
19. ABSTRACT (Continue on reverse side if necessary and identify by block number) Stochastic dominance orders of all finite degrees are defined on the set of distribution functions on the nonnegative real numbers in terms of integrals of the distributions. It is proved that if F strictly nth-degree stochastically dominates G, and if the moments of F and G through order n are finite with $\mu_F^k = \int x^k dF(x)$ , then $(\mu_F^1, \dots, \mu_F^n) \neq (\mu_G^1, \dots, \mu_G^n)$ and $(-1)^{k-1} \mu_F^k > (-1)^{k-1} \mu_G^k$ for the smallest k for which $\mu_F^k \neq \mu_G^k$ .		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED


SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

407732

78 06 26 047

AD IN. DDC FILE COPY

AD A055829



DDC		Diff Section	<input type="checkbox"/>
UNANNOUNCED			<input checked="" type="checkbox"/>
JUSTIFICATION			
BY			
DISTRIBUTION/AVAILABILITY CODES			
DISTRIBUTION/AVAILABILITY CODES			
SPECIAL			
A			

# TECHNICAL REPORT NO. 29

## STOCHASTIC DOMINANCE AND MOMENTS OF DISTRIBUTIONS

Peter C. Fishburn  
 College of Business Administration  
 The Pennsylvania State University  
 University Park, Pennsylvania 16802

This research was sponsored by the Office of Naval Research, ONR Contract Number N00014-75-C-0857, ONR Contract Authority identification number NR047-112/11-4-77.

Reproduction in whole or in part is permitted for any purpose of the United States Government.

78 06 26 047

## 1. Introduction

Stochastic dominance provides a way of analyzing risky decisions when a decision agent's von Neumann-Morgenstern (1947) utility function  $u$  is not fully known but is presumed to be in a class  $U$  of real-valued functions defined on a consequence space  $X$ . When  $p$  and  $q$  are probability measures on a suitable algebra of subsets of  $X$ , we sometimes say that  $p$  stochastically dominates  $q$  with respect to  $U$  if  $\int u(x)dp(x) > \int u(x)dq(x)$  for all  $u \in U$  for which the expected utilities are finite.

Although stochastic dominance is relevant whenever decision alternatives are described by probability measures, its most popular use involves situations in which  $X$  is a set of real numbers with  $x$  preferred to  $y$  when  $x > y$ . This context is emphasized in the introduction to stochastic dominance given by Fishburn and Vickson (1978) and will be the context used for the present paper. Definitions of various degrees of stochastic dominance in the real  $X$  setting can be given without explicit mention of  $U$  classes, and I will follow this approach here.

My purpose is to establish a general connection between  $n$ th-degree stochastic dominance and the first  $n$  moments of probability measures whose supports are bounded below, for all  $n \in \{1, 2, \dots\}$ . It will be assumed that the measures are countably additive and that their supports are bounded below by 0. The corresponding distribution functions  $F, G, \dots$  will be taken to be continuous from the right with  $F(x) = 0$  for all  $x < 0$  and  $F(x) \rightarrow 1$  as  $x \rightarrow \infty$ .

The basic definitions and main theorem are presented and discussed in the next section where the main theorem is observed to follow from two auxiliary theorems. Proofs of the latter theorems are given in the final two sections. These proofs are based solely on the distribution functions and will not involve utility classes.



## 2. Definitions and Theorems

Let  $F$  be the set of all right-continuous distribution functions on the real line with  $F(x) = 0$  for all  $x < 0$  for each  $F \in F$ . For each  $F \in F$  let  $F^1 = F$  and recursively define

$$F^{n+1}(x) = \int_0^x F^n(y) dy \quad \text{for all } x \geq 0 \text{ and } n \in \{1, 2, \dots\}.$$

Nonstrict ( $\geq_n$ ) and strict ( $>_n$ )  $n$ th-degree stochastic dominance relations are then defined on  $F$  as follows:

$$F \geq_n G \text{ iff } F^n(x) \leq G^n(x) \quad \text{for all } x \in [0, \infty),$$

$$F >_n G \text{ iff } F \neq G \text{ and } F \geq_n G.$$

These relations are transitive and increasingly more inclusive:  $\geq_1 \subset \geq_2 \subset \dots$  and  $>_1 \subset >_2 \subset \dots$  where  $\subset$  denotes proper inclusion. Utility classes that are congruent with the first few stochastic dominance orders are presented in Fishburn and Vickson (1978, pp. 102-113). For example,  $F \geq_1 G$  iff  $\int u(x) dF(x) \geq \int u(x) dG(x)$  for all nondecreasing  $u$  on  $[0, \infty)$  for which the expectations exist, and  $F \geq_2 G$  iff  $\int u dF \geq \int u dG$  for all nondecreasing and concave  $u$  on  $[0, \infty)$  for which the expectations exist. The corresponding theorems for  $(>_1, >)$  and  $(>_2, >)$  respectively involve strictly increasing  $u$  and strictly increasing-strictly concave  $u$ . Congruent utility classes for the third-degree relations are the subsets of the second-degree classes in which first derivatives exist and are convex ( $\geq_3$ ) or strictly convex ( $>_3$ ). Although it is traditional to define  $F \geq_3 G$  iff  $F^3(x) \leq G^3(x)$  for all  $x$  and the mean of  $F$  is as great as the mean of  $G$  (e.g., [Whitmore, 1970], [Fishburn and Vickson, 1978]), we shall see that the condition on the means is redundant in the present formulation.

The moment sequence for  $F \in \mathcal{F}$  is  $\mu(F) = (\mu_F^0, \mu_F^1, \mu_F^2, \dots)$  where  $\mu_F^0 = \int dF(x) = 1$  and  $\mu_F^n = \int x^n dF(x)$  for each  $n \geq 1$ . The part of this sequence through the  $n$ th moment will be denoted as  $\mu_n(F) = (\mu_F^0, \dots, \mu_F^n)$ . Our main theorem will be based on binary relations  $>_n^*$  on  $\{\mu(F): F \in \mathcal{F}\}$  defined lexicographically on the basis of  $\mu_n$  as follows:

$$\mu(F) >_n^* \mu(G) \text{ iff } \mu_n(F) \neq \mu_n(G) \text{ and } (-1)^{k-1} \mu_F^k > (-1)^{k-1} \mu_G^k$$

for the smallest  $k$  for which  $\mu_F^k \neq \mu_G^k$ .

For example,  $\mu(F) >_1^* \mu(G)$  iff  $\mu_F^1 > \mu_G^1$ , and  $\mu(F) >_2^* \mu(G)$  iff either  $\mu_F^1 > \mu_G^1$  or  $(\mu_F^1 = \mu_G^1, \mu_F^2 < \mu_G^2)$ . Alternatively,  $\mu(F) >_2^* \mu(G)$  iff either  $F$  has the greater mean or the means of  $F$  and  $G$  are equal and  $F$  has a smaller variance than  $G$ .

**Theorem 1.** If  $F, G \in \mathcal{F}$  and the moments of  $F$  and  $G$  through order  $n$  are finite, then  $\mu(F) >_n^* \mu(G)$  if  $F >_n G$ .

For discussion purposes the next three paragraphs assume that the moments involved therein are finite.

The  $n = 1$  part of Theorem 1 says that if  $F >_1 G$  then the mean of  $F$  must be greater than the mean of  $G$ . Equivalently, if  $\mu_F^1 \leq \mu_G^1$  then  $F$  cannot strictly first-degree stochastically dominate  $G$ . This fact seems to be widely known.

It appears to be less commonly recognized that  $F >_2 G$  implies that either  $\mu_F^1 > \mu_G^1$  (in which case the inequality on second moments could go either way) or  $(\mu_F^1 = \mu_G^1, \mu_F^2 < \mu_G^2)$ . Parts of this result have been noted by Hanoch and Levy (1969), Rothschild and Stiglitz (1970) and Fishburn and Vickson (1978, p. 78).

For  $n = 3$ , Theorem 1 says that  $F >_3 G$  entails either  $\mu_F^1 > \mu_G^1$  or  $(\mu_F^1 = \mu_G^1, \mu_F^2 < \mu_G^2)$  or  $(\mu_F^1 = \mu_G^1, \mu_F^2 = \mu_G^2, \mu_F^3 > \mu_G^3)$ . Although parts of this have been noted by Whitmore (1970) and Fishburn and Vickson (1978, pp. 78-82) for special cases, the complete result appears to be previously unknown. In like manner, Theorem 1 for  $n \geq 4$  appears to be new.

Since  $F >_3 G$  implies  $\mu_F^1 \geq \mu_G^1$  when these means are finite--and even when one or both are infinite--our definition of  $>_3$  (or  $\geq_3$ ) is formally equivalent to the traditional definition. It should be noted however that upper-bounded definitions are not so simply related. In particular, if  $[0, b]$  with  $b$  finite includes the supports of  $F$  and  $G$  then, as shown in Fishburn (1976), it is possible to have  $\mu_F^1 < \mu_G^1$  when  $F^3(x) \leq G^3(x)$  for all  $x \in [0, b]$ . In this case, when  $F^3(x) \leq G^3(x)$  is specified only over an interval that includes the supports of  $F$  and  $G$ , it is necessary to include the stipulation  $\mu_F^1 \geq \mu_G^1$  in the definition of  $F \geq_3 G$  if we want  $\geq_3$  to be congruent with the type of  $U$  class mentioned in the opening paragraph of this section.

The proof of Theorem 1 can be based on the following two auxiliary theorems that will be proved in the next two sections.

Theorem 2. If  $F, G \in \mathcal{F}$ , if  $F \geq_m G$  for some  $m \in \{1, 2, \dots\}$ , and if the moments of  $F$  and  $G$  through order  $n + 1 \geq 1$  are finite with  $\mu_F^k = \mu_G^k$  for  $k = 0, 1, \dots, n$ , then  $(-1)^n \mu_F^{n+1} \geq (-1)^n \mu_G^{n+1}$ .

Theorem 3. If  $F, G \in \mathcal{F}$ , if  $F >_n G$ , and if the moments of  $F$  and  $G$  through order  $n \geq 1$  are finite with  $\mu_F^k = \mu_G^k$  for  $k = 0, 1, \dots, n - 1$ , then  $(-1)^{n-1} \mu_F^n > (-1)^{n-1} \mu_G^n$ .

The latter theorem implies that  $\mu_n(F) = \mu_n(G)$  cannot be true when  $F >_n G$ , and the former theorem says that if  $F$  stochastically dominates  $G$  in the



nonstrict sense for any finite degree and if  $\mu_n(F) = \mu_n(G)$  then  $\mu_F^{n+1} \geq \mu_G^{n+1}$  if  $n$  is even and  $\mu_F^{n+1} \leq \mu_G^{n+1}$  if  $n$  is odd, provided that the moments involved are finite. Theorem 1 follows immediately from Theorems 2 and 3: if  $F >_n G$  then  $F \geq_n G$ , hence either  $\mu(F) >^*_n \mu(G)$  or  $\mu_n(F) = \mu_n(G)$  by Theorem 2; since Theorem 3 rules out  $\mu_n(F) = \mu_n(G)$  we are left with  $\mu(F) >^*_n \mu(G)$ .

### 3. Proof of Theorem 2

We begin with a lemma that leads to the proper sense of the inequality in the conclusion of Theorem 2.

Lemma 1. For any  $H \in F$  with finite moments through order  $n$  and for all  $m > n \geq 0$  and  $x > 0$  let

$$T_{n,m}(H,x) = \sum_{k=0}^n (-1)^{k+1} \binom{m}{k} \mu_H^k / x^k + \int_0^x (1 - y/x)^m dH(y). \quad (1)$$

Then  $T_{n,m}(H,x) = 0$  iff  $H(0) = 1$  and, for all other  $H \in F$ ,  $T_{n,m}(H,x) > 0$  if  $n$  is odd, and  $T_{n,m}(H,x) < 0$  if  $n$  is even.

Proof. If  $H$  assigns probability mass 1 to  $y = 0$  then (1) gives  $T_{n,m}(H,x) = -1 + 1 = 0$ . Assume henceforth that  $H(0) < 1$ . When  $\mu_H^k$  in the right hand side of (1) is replaced by  $\int_0^x y^k dH(y) + \int_x^\infty y^k dH(y)$ , and  $(1 - y/x)^m$  is expanded binomially, we get

$$\begin{aligned} T_{n,m}(H,x) &= \int_0^x \left\{ \sum_{k=n+1}^m (-1)^k \binom{m}{k} (y/x)^k \right\} dH(y) + \int_x^\infty \left\{ \sum_{k=0}^n (-1)^{k+1} \binom{m}{k} (y/x)^k \right\} dH(y) \\ &= \int_0^x A_{n,m}(z) dH(y) + \int_x^\infty B_{n,m}(z) dH(y) \end{aligned} \quad (2)$$



where the A and B terms are defined in context as the sums under the integrals and  $z = y/x$  with  $z \in [0,1]$  for A and  $z \in [1,\infty)$  for B. We consider A and B in turn.

First,  $A_{n,m}(0) = 0$  for all  $m > n \geq 0$  and, since  $A_{0,m}(z) = (1-z)^m - 1$ ,  $A_{0,m}(z) < 0$  for all  $z \in (0,1]$ . Since

$$A'_{n,m}(z) = dA_{n,m}(z)/dz = -mA_{n-1,m-1}(z) \text{ for } m > n \geq 1 \text{ and } z > 0,$$

it follows that  $A'_{1,m}(z) > 0$  for  $z > 0$ , hence by continuity and  $A_{1,m}(0) = 0$  that  $A_{1,m}(z) > 0$  for all  $z \in (0,1]$  and  $m > 1$ . Then  $A'_{2,m}(z) < 0$  for  $z > 0$  and  $m > 2$ , so that  $A_{2,m}(z) < 0$  for all  $z \in (0,1]$ . The obvious continuation of this process gives

$$A_{n,m}(z) > 0 \text{ for all } z \in (0,1] \text{ if } n \text{ is odd,}$$

$$A_{n,m}(z) < 0 \text{ for all } z \in (0,1] \text{ if } n \text{ is even,}$$

along with  $A_{n,m}(0) = 0$ .

For the B part we note first (e.g. [Feller, 1957, p. 61]) that  $B_{n,m}(1) = (-1)^{n+1} \binom{m-1}{n}$  so that  $B_{n,m}(1) > 0$  if  $n$  is odd, and  $B_{n,m}(1) < 0$  if  $n$  is even. In addition,

$$B'_{n,m}(z) = -m B_{n-1,m-1}(z) \text{ for } m > n \geq 1 \text{ and } z \geq 1.$$

Since  $B_{0,m}(z) \equiv -1$ , this implies that  $B'_{1,m}(z) > 0$  for  $m > 1$  and  $z \geq 1$ , hence that  $B_{1,m}(z) > 0$  for all  $z \geq 1$  since  $B_{1,m}(1) > 0$ . Then  $B'_{2,m}(z) < 0$ , hence  $B_{2,m}(z) < 0$  for all  $z \geq 1$  since  $B_{2,m}(1) < 0$ . The continuation of this process shows that

$B_{n,m}(z) > 0$  for all  $z \in [1, \infty)$  if  $n$  is odd,

$B_{n,m}(z) < 0$  for all  $z \in [1, \infty)$  if  $n$  is even.

These conclusions along with those for  $A$  and the hypothesis that  $H$  has positive probability mass on  $x > 0$  then yield the final conclusions of Lemma 1 in view of (2). Q.E.D.

For Theorem 2, suppose that  $F, G \in \mathcal{F}$  have finite moments through order  $n+1 \geq 1$  and that  $\mu_n(F) = \mu_n(G)$ , i.e.  $\mu_F^k = \mu_G^k$  for all  $k \leq n$ . Assume also that  $F \geq_m G$  for some  $m$ . Since  $\geq_m \subset \geq_{m+1}$  for all  $m$ , select  $m > n$  with  $F \geq_{m+1} G$ . Then, by the definition of  $\geq_{m+1}$  and Fishburn (1976),

$$\int_0^x (x-y)^m dF(y) \leq \int_0^x (x-y)^m dG(y) \text{ for all } x > 0.$$

Since  $\mu_n(F) = \mu_n(G)$ , it follows that

$$T_{n,m}(G, x) \geq T_{n,m}(F, x) \text{ for all } x > 0. \quad (3)$$

Suppose first that  $G(0) = 1$  so that  $T_{n,m}(G, x) = 0$  by Lemma 1. Then  $0 \geq T_{n,m}(F, x)$  for all  $x > 0$ . It follows that either  $F(0) = 1$  and  $F = G$ , in which case  $\mu_F^{n+1} = \mu_G^{n+1} = 0$ , or that  $F(0) < 1$ , in which case Lemma 1 implies that  $0 > T_{n,m}(F, x)$  for  $x > 0$  and hence that  $n$  is even. But with  $n$  even,  $(-1)^n \mu_F^{n+1} > (-1)^n \mu_G^{n+1} = 0$  in the latter case. Hence the conclusion of Theorem 2 holds if  $G(0) = 1$ .

Assume henceforth that  $G(0) < 1$ , in which case  $F(0) < 1$  also. Then (3) applies with neither term ever equal to zero according to the last part of Lemma 1. We shall prove that

$$\lim_{x \rightarrow \infty} \frac{T_{n,m}(F, x)}{T_{n,m}(G, x)} = \frac{\mu_F^{n+1}}{\mu_G^{n+1}} \quad (4)$$

which, in view of (3) and the final conclusions of the lemma, implies that  $(-1)^n \mu_F^{n+1} \geq (-1)^n \mu_G^{n+1}$ , as desired for Theorem 2. Since  $\lim_{n,m} T_{n,m}(H,x) = -\mu_H^0 + 1 = 0$ , we differentiate the numerator and denominator of the left hand side of (4) with respect to  $x$  in anticipation of using l'Hospital's rule. When  $m/x^2$  is canceled from the numerator and denominator after the differentiations have been completed, we get

$$\frac{d T_{n,m}(F,x)/dx}{d T_{n,m}(G,x)/dx} = \frac{\sum_{k=0}^{n-1} (-1)^{k+1} \binom{m-1}{k} \mu_F^{k+1}/x^k + \int_0^x (1-y/x)^{m-1} y dF(y)}{\sum_{k=0}^{n-1} (-1)^{k+1} \binom{m-1}{k} \mu_G^{k+1}/x^k + \int_0^x (1-y/x)^{m-1} y dG(y)} . \quad (5)$$

If  $n = 0$  then the numerator and denominator of (5) respectively approach  $\mu_F^1$  and  $\mu_G^1$  as  $x \rightarrow \infty$  by the monotone convergence theorem (e.g., [Loeve, 1960, p. 124]), and a single application of l'Hospital's rule shows that (4) holds. If  $n > 1$ , then the numerator and denominator of the right hand side of (5) both go to zero ( $-\mu_F^1 + \mu_F^1$  and  $-\mu_G^1 + \mu_G^1$ ) as  $n \rightarrow \infty$ . In this case methods similar to those used for Lemma 1 show that both parts of the right hand side of (5) have constant nonzero sign. We then differentiate the numerator and denominator of the right hand side of (5) and cancel  $(m-1)/x^2$  to get

$$\frac{\sum_{k=0}^{n-2} (-1)^{k+1} \binom{m-2}{k} \mu_F^{k+2}/x^k + \int_0^x (1-y/x)^{m-2} y^2 dF(y)}{\sum_{k=0}^{n-2} (-1)^{k+1} \binom{m-2}{k} \mu_G^{k+2}/x^k + \int_0^x (1-y/x)^{m-2} y^2 dG(y)} .$$

If  $n = 1$  then (4) holds by two applications of l'Hospital's rule since the limit of the expression just written is  $\mu_F^2/\mu_G^2$ . If  $n > 1$ , we continue in the indicated



manner until we arrive at the ratio  $\int_0^x (1 - y/x)^{m-n-1} y^{n+1} dF(y) / \int_0^x (1 - y/x)^{m-n-1} y^{n+1} dG(y)$ , whose limit is  $\mu_F^{n+1} / \mu_G^{n+1}$ . Hence  $n + 1$  applications of l'Hospital's rule yield a general verification of (4).

#### 4. Proof of Theorem 3

Throughout this section we use the convention that  $t^0 = 1$  when  $t = 0$ . In anticipation of using the monotone convergence theorem later we shall first prove

**Lemma 2.** Suppose  $H \in F$  has finite moments through order  $n - 1$  with  $n \geq 1$ , and  $0 \leq y \leq x$  with  $x > 0$ . Then

$$\begin{aligned} & (-1)^n [n! H^n(y) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} k \mu_H^{n-k} y^{k-1}] \\ & \geq (-1)^n [n! H^n(y) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} k y^{k-1} \int_{z=0}^x z^{n-k} dH(z)] \\ & \geq 0. \end{aligned}$$

**Proof.** We note first that

$$\begin{aligned} \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} k \mu_H^{n-k} y^{k-1} &= n \sum_{k=0}^{n-1} (-1)^{n-k} \binom{n-1}{k} y^k \mu_H^{n-k-1}, \\ \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} k y^{k-1} \int_{z=0}^x z^{n-k} dH(z) &= n \sum_{k=0}^{n-1} (-1)^{n-k} \binom{n-1}{k} y^k \int_{z=0}^x z^{n-k-1} dH(z) \\ &= -n \int_{z=0}^x (y - z)^{n-1} dH(z), \\ n! H^n(y) &= n \int_{z=0}^y (y - z)^{n-1} dH(z) \quad ([Fishburn, 1976]). \end{aligned}$$

By the first two of these, the first inequality in the conclusion of Lemma 2 holds if and only if

$$n(-1)^n \left[ \sum_{k=0}^{n-1} (-1)^{n-k} \binom{n-1}{k} y^k \left\{ \mu_H^{n-k-1} - \int_{z=0}^x z^{n-k-1} dH(z) \right\} \right] \geq 0,$$

which is true if and only if

$$\sum_{k=0}^{n-1} (-1)^k \binom{n-1}{k} y^k \int_{z=x}^{\infty} z^{n-k-1} dH(z) \geq 0.$$

But this is true since its left hand side equals  $\int_{z=x}^{\infty} (z-y)^{n-1} dH(z)$ . By the final two expressions in the opening sentence of this proof, the second ( $\geq 0$ ) inequality in Lemma 2 holds if

$$(-1)^n \left[ -n \int_{z=y}^x (y-z)^{n-1} dH(z) \right] \geq 0,$$

which is true since its left hand side equals  $n \int_{z=y}^x (z-y)^{n-1} dH(z)$ . Q.E.D.

When  $dy$  is replaced by  $e^y d(1 - e^{-y})$  in the following expressions, with  $1 - e^{-y}$  a distribution function in  $F$ , Lemma 2 in conjunction with the monotone convergence theorem shows that if  $\mu_H^{n-1}$  is finite then

$$\begin{aligned} & (-1)^n \int_{y=0}^{\infty} [n! H^n(y) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} \mu_H^{n-k} y^{k-1}] dy \\ &= \lim_{x \rightarrow \infty} \{ (-1)^n \int_{y=0}^x [n! H^n(y) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} y^{k-1} \int_{z=0}^x z^{n-k} dH(z)] dy \}. \end{aligned}$$

The expression within braces after  $\lim$  reduces as follows:

$$\begin{aligned} & (-1)^n [n! H^{n+1}(x) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} x^k \int_{z=0}^x z^{n-k} dH(z)] \\ &= (-1)^n \left[ \int_{y=0}^x (x-y)^n dH(y) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} x^k \int_{y=0}^x y^{n-k} dH(y) \right] \end{aligned}$$

$$\begin{aligned}
&= (-1)^n \left[ \int_{y=0}^x \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} x^k y^{n-k} dH(y) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} x^k \int_{y=0}^x y^{n-k} dH(y) \right] \\
&= \int_{y=0}^x y^n dH(y).
\end{aligned}$$

Therefore, when  $\mu_H^{n-1}$  is finite,

$$\int_{y=0}^{\infty} [n! H^n(y) + \sum_{k=1}^n (-1)^{n-k+1} \binom{n}{k} k \mu_H^{n-k} y^{k-1}] dy = (-1)^n \mu_H^n. \quad (6)$$

Finally, as in the hypotheses of Theorem 3, suppose that  $F >_n G$  and that the moments of  $F$  and  $G$  through order  $n$  are finite with  $\mu_F^k = \mu_G^k$  for all  $k \leq n-1$ . Then, since  $F^n(y) \leq G^n(y)$  for all  $y$ , and  $F^n(y) < G^n(y)$  in a non-degenerate interval within  $[0, \infty)$ , (6) shows immediately that  $(-1)^n \mu_F^n < (-1)^n \mu_G^n$ .



## References

- [1] Feller, W. (1957). An Introduction to Probability Theory and Its Applications. 2nd Ed., Wiley, New York.
- [2] Fishburn, P. C. (1976). Continua of Stochastic Dominance Relations for Bounded Probability Distributions. J. Math. Econ. 3 295-311.
- [3] \_\_\_\_\_ and Vickson, R. G. (1978). Theoretical Foundations of Stochastic Dominance. In Stochastic Dominance (G. A. Whitmore and M. C. Findlay, eds.). D. C. Heath and Co., Lexington, Massachusetts, 37-113.
- [4] Hanoch, G. and Levy, H. (1969). The Efficiency Analysis of Choices Involving Risk. Rev. Econ. Studies 36 335-346.
- [5] Loève, M. (1960). Probability Theory. 2nd Ed., Van Nostrand, Princeton, New Jersey.
- [6] Rothschild, M. and Stiglitz, J. E. (1970). Increasing Risk: I. A Definition. J. Econ. Theory 2 225-243.
- [7] von Neumann, J. and Morgenstern, O. (1947). Theory of Games and Economic Behavior. 2nd Ed., Princeton University Press, Princeton, New Jersey.
- [8] Whitmore, G. A. (1970). Third-Degree Stochastic Dominance. Amer. Econ. Rev. 60 457-459.

1 Oct. 78

BASIC DISTRIBUTION LIST FOR  
UNCLASSIFIED TECHNICAL REPORTS  
Operations Research Program

Code 434

Operations Research Office of Naval Research (Code 434) Arlington, Virginia 22217	3	Library Naval Electronics Laboratory Center San Diego, California 92152	1
Director, Naval Research Laboratory Attn: Library, Code 2029 (ONRL) Washington, D.C. 20390	2	Dr. A. L. Slafkosky Scientific Advisor Commandant Marine Corps (Code AX) Washington, D.C. 20380	1
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	12	Naval Undersea Warfare Center 3202 E. Foothill Boulevard Pasadena, California 91107 Attn: Technical Library	1
Defense Logistics Studies Information Exchange Army Logistics Management Center Fort Lee, Virginia 23801	1	Stanford University Department of Statistics Stanford, California 94305 Attn: Prof. G. J. Lieberman	1
Technical Information Center Naval Research Laboratory Washington, D.C. 20390	6	Stanford University Department of Statistics Stanford, California 94305 Attn: Prof. T. W. Anderson	1
Director, Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91106 Attn: Dr. R. Lau	1	University of California Department of Engineering Los Angeles, California 90024 Attn: Prof. R. R. O'Neill	1
Mr. William Dejka Naval Electronics Laboratory Center, Code 360 San Diego, California 92132	1	Maritime Transportation Research Board National Academy of Sciences 2101 Constitution Avenue Washington, D.C. 20418 Attn: RADM J. B. Oren, USCG (RET)	1
Director Office of Naval Research Branch Office 495 Summer Street Boston, Massachusetts 02210 Attn: Dr. A. L. Powell	1	Director Office of Naval Research Branch Office 536 South Clark Street Chicago, Illinois 60605 Attn: Dr. R. Buchal	1
Library Naval Postgraduate School Code 0212 Monterey, California 93940	1	Stanford University Department of Operations Research Stanford, California 94305 Attn: Prof. A. F. Veinott, Jr.	1
Naval Ordnance Station Louisville, Kentucky 40214	1		

Texas A&M Foundation College Station, Texas 77843 Attn: Prof. H. O. Hartley	1	University of Texas Department of Mathematics Austin, Texas 78712 Attn: Dr. A. Charnes	1
U.S. Naval Postgraduate School Department of Operations Research and Administrative Services Monterey, California 93940	1	Stanford University Department of Operations Research Stanford, California 94305 Attn: Prof. F. S. Hillier	1
Captain Robert Bush AFOSR (SRMA) 1400 Wilson Blvd. Arlington, Virginia 22209	1	Stanford University Department of Operations Research Stanford, California 94305 Attn: Prof. G. B. Dantzig	1
Professor Richard L. Fox School of Engineering Case Western Reserve University Cleveland, Ohio 44106	1	Cornell University Department of Industrial Engineering Ithaca, New York 19850 Attn: Prof. W. Lucas	1
Director, National Security Agency Ft. George G. Meade, Maryland 20755 Attn: Dr. Joseph Blum, R44	1	Dr. Jay Lewallen E-D-13 Theory and Analysis Division NASA Manned Spacecraft Center Houston, Texas 77058	1
Logistics Research Project The George Washington University 707 22nd Street, N.W. Washington, D.C. 20037 Attn: Dr. W. H. Marlow	1	Applied Mathematics Laboratory Naval Ships Research Development Center Washington, D.C. 20007	1
Harvard University Department of Economics Cambridge, Massachusetts 02138 Attn: Prof. K. J. Arrow	1	Mr. A. S. Rhode Office of Chief of Naval Operations Code 964 Arlington, Virginia 22203	2
Cowles Commission for Research in Economics Yale University New Haven, Connecticut 06520 Attn: Prof. Martin Shubik	1	Dr. Del Cilmer Naval Weapons Center (Code 607) China Lake, California 93555	1
University of California Operations Research Center Institute of Engineering Research Berkeley, California 94720 Attn: Prof. D. Gale	1	Carnegie-Mellon University Graduate School of Industrial Adm. Pittsburgh, Pennsylvania 15213 Attn: Prof. G. Thompson	1
The University of Michigan Department of Mathematics, W.E. Ann Arbor, Michigan 48104 Attn: Prof. R. M. Thrall	1	Dr. Thom J. Hodgson Department of Industrial and Systems Engineering University of Florida Gainesville, Florida 32601	1



Professor Salah E. Elmaghraby  
Department of Operations Research  
and Industrial Engineering  
North Carolina State University  
Raleigh, North Carolina 27607

1

Mr. Thaddeus G. Bell  
Navy Underwater Sound Laboratory  
Fort Trumbull  
New London, Connecticut 06320

1

Dr. Jack R. Borsting, Chairman  
Dept. of Operations Research and  
Administrative Sciences  
Naval Postgraduate School  
Monterey, California 93940

1

Dr. G. S. Colladay  
Naval Undersea Research  
and Development Center  
San Diego, California 92132

1

Professor F. J. Gould  
Graduate School of Business  
University of Chicago  
5836 Greenwood Avenue  
Chicago, Illinois 60637

1

Professor Joseph L. Balintfy  
Department of General Business  
and Finance  
School of Business Administration  
University of Massachusetts  
Amherst, Massachusetts 01002

1

Professor H. Donald Ratliff  
Department of Industrial and  
Systems Engineering  
University of Florida  
Gainesville, Florida 32601

1

Prof. Leon Lasdon  
Operations Research Dept.  
Case Western Reserve University  
Cleveland, Ohio 44106

1

Professor A. Geoffrion  
Graduate School of Business Adm.  
University of California  
Los Angeles, California 90024

1

Dr. Peter C. Fishburn  
Research Professor of Management  
Science  
College of Business Administration  
The Pennsylvania State University  
University Park, Pa. 16802

1

Professor Ralph L. Keeney  
Woodward-Clyde Consultants  
Three Embarcadero Center  
Suite 700  
San Francisco, California 94111

1

Assistant Chief for Technology  
Office of Naval Research, Code 200  
Arlington, Virginia 22217

1